

CORRECTION

Open Access



Correction to: Type 1 diabetes susceptibility alleles are associated with distinct alterations in the gut microbiota

Jane A. Mullaney^{1,2}, Juliette E. Stephens¹, Mary-Ellen Costello³, Cai Fong¹, Brooke E. Geeling¹, Patrick G. Gavin¹, Casey M. Wright¹, Timothy D. Spector⁴, Matthew A. Brown³ and Emma E. Hamilton-Williams^{1,5*}

Correction

Following publication of the original article [1] it came to the attention of the Production Editor that Figs. 1 and 2 had not been replaced with the newly revised figures supplied by the authors (the originals being unusable due to poor quality image and text).

The original article has been corrected.

Author details

¹Translational Research Institute, The University of Queensland Diamantina Institute, University of Queensland, QLD, Brisbane, Australia. ²AgResearch Limited, Grasslands Research Centre, Palmerston North, New Zealand.

³Institute of Health and Biomedical Innovation, Queensland University of Technology, Translational Research Institute, QLD, Brisbane, Australia.

⁴Department of Twin Research and Genetic Epidemiology, King's College

London, London SE1 7EH, UK. ⁵Translational Research Institute, The University of Queensland Diamantina Institute, 37 Kent St, Woolloongabba, QLD 4102, Australia.

Received: 12 March 2018 Accepted: 12 March 2018

Published online: 20 March 2018

Reference

1. Mullaney JA, Stephens JE, Costello ME, Fong C, Geeling BE, Gavin PG, Wright CM, Spector TD, Brown MA, Hamilton-Williams EE. Type 1 diabetes susceptibility alleles are associated with distinct alterations in the gut microbiota. *Microbiome*. 2018;6:35. <https://doi.org/10.1186/s40168-018-0417-4>.

* Correspondence: e.hamiltonwilliams@uq.edu.au

¹Translational Research Institute, The University of Queensland Diamantina Institute, University of Queensland, QLD, Brisbane, Australia

⁵Translational Research Institute, The University of Queensland Diamantina Institute, 37 Kent St, Woolloongabba, QLD 4102, Australia