## CORRECTION Open Access



## Correction: Darier's disease exhibits a unique cutaneous microbial dysbiosis associated with inflammation and body malodour

Yacine Amar<sup>1†</sup>, Danielle Rogner<sup>1†</sup>, Rafaela L. Silva<sup>1</sup>, Bärbel U. Foesel<sup>2</sup>, Minhaz Ud-Dean<sup>3,4</sup>, Ilias Lagkouvardos<sup>5</sup>, Susanne A. Steimle-Grauer<sup>1</sup>, Sebastian Niedermeier<sup>1</sup>, Susanne Kublik<sup>2</sup>, Manja Jargosch<sup>1</sup>, Matthias Heinig<sup>3,4</sup>, Jenny Thomas<sup>6</sup>, Stefanie Eyerich<sup>6</sup>, Jakob D. Wikström<sup>7,8</sup>, Michael Schloter<sup>2</sup>, Kilian Eyerich<sup>7,9</sup>, Tilo Biedermann<sup>1\*</sup> and Martin Köberle<sup>1</sup>

Correction: Microbiome 11, 162 (2023) https://doi.org/10.1186/s40168-023-01587-x

Following publication of the original article [1], the author reported that the name "Jakob Wikström" should be "Jakob D Wikström"

This has been corrected above and the original article has been updated.

<sup>†</sup>Yacine Amar and Danielle Rogner equally contributed to this study.

The original article can be found online at https://doi.org/10.1186/s40168-023-01587-x.

\*Correspondence:

Tilo Biedermann

tilo.biedermann@tum.de

- <sup>1</sup> Department of Dermatology and Allergy, Technical University of Munich, School of Medicine, Munich, Germany
- <sup>2</sup> Research Unit Comparative Microbiome Analysis, Helmholtz Zentrum München, Deutsches Forschungszentrum Für Gesundheit Und Umwelt (GmbH), 85764 Neuherberg, Germany
- <sup>3</sup> Institute of Computational Biology, Helmholtz Zentrum München, German Research Center for Environmental Health, 85764 Neuherberg, Germany
- <sup>4</sup> Department of Informatics, Technical University of Munich, Garching, Germany
- <sup>5</sup> Core Facility Microbiome, Technical University of Munich, 85354 Freising, Germany
- <sup>6</sup> Center of Allergy & Environment (ZAUM), Technical University of Munich (TUM) and Helmholtz Zentrum München, German Research Center for Environmental Health, Member of the German Center of Lung Research (DZL). Munich. Germany
- Dermatology and Venereology Division, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden

Published online: 14 September 2023

## Reference

 Amar Y, Rogner D, Silva RL, et al. Darier's disease exhibits a unique cutaneous microbial dysbiosis associated with inflammation and body malodour. Microbiome. 2023;11:162. https://doi.org/10.1186/ s40168-023-01587-x.



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data

<sup>&</sup>lt;sup>8</sup> Dermato-Venereology Clinic, Karolinska University Hospital, Stockholm, Sweden

<sup>&</sup>lt;sup>9</sup> Department of Dermatology and Venereology, Medical Center, University of Freiburg, Freiburg, Germany