CORRECTION Open Access

Correction to: Metagenomic analysis of isolation methods of a targeted microbe, *Campylobacter jejuni*, from chicken feces with high microbial contamination



Junhyung Kim¹, Jae-Ho Guk¹, Seung-Hyun Mun¹, Jae-Uk An¹, Hyokeun Song¹, Jinshil Kim², Sangryeol Ryu², Byeonghwa Jeon^{2,3} and Seongbeom Cho^{1*}

Correction to: Microbiome https://doi.org/10.1186/s40168-019-0680-z

Following publication of the original article [1], the authors reported an error in Fig. 2. The correct figure is shown below.

Author details

¹Research Institute for Veterinary Science and College of Veterinary Medicine, Seoul National University, Seoul, Republic of Korea. ²Department of Food and Animal Biotechnology, Department of Agricultural Biotechnology, Research Institute for Agriculture and Life Sciences, and Center for Food and Bioconvergence, Seoul National University, Seoul, Republic of Korea. ³School of Public Health, University of Alberta, Edmonton, Alberta, Canada.

Published online: 01 June 2019

Reference

 Kim J, Guk J-H, Mun S-H, An J-U, Song H, Kim J, Ryu S, Jeon B, Cho S. Metagenomic analysis of isolation methods of a targeted microbe, Campylobacter jejuni, from chicken feces with high microbial contamination. Microbiome. 2019;7:67.

¹Research Institute for Veterinary Science and College of Veterinary Medicine, Seoul National University, Seoul, Republic of Korea Full list of author information is available at the end of the article



^{*} Correspondence: chose@snu.ac.kr

Kim et al. Microbiome (2019) 7:81 Page 2 of 2




Fig. 2 Proportion of competing microbes according to isolation procedures of *C. jejuni* based on culture-dependent tools. Proportion of **a** ESBL-producing *E. coli* and **b** *P. mirabilis* among competing microbes according to the isolation procedure of *C. jejuni* (mean \pm SEM). Significance was determined by *t* test and one-way ANOVA. The proportion of ESBL-producing *E. coli* was significantly different according to the isolation procedure including the type of enrichment broth, selective media, and the combination of different enrichment broths, the ratio of sample-to-enrichment broth, and selective media. The proportion of *P. mirabilis* was significantly different according to the isolation procedure including the type of selective media and combination of different enrichment broths, the ratio of sample to enrichment broth, and selective agars. *p < 0.05, **p < 0.05 (*p <